

Cardiorespiratory Endurance Assessment of High School Futsal Players: A Case Study at SMAS Fatih Bilingual School, Banda Aceh

by Fachrul Azmi

Submission date: 14-Aug-2025 08:08PM (UTC+0530)

Submission ID: 2729555760

File name: PUBLISH_AZMI_IJESS_13-16.pdf (503.85K)

Word count: 2253

Character count: 13639



Cardiorespiratory Endurance Assessment of High School Futsal Players: A Case Study at SMAS Fatih Bilingual School, Banda Aceh

Fachrul Azmi*, Ifwandi, Masri, Zulfikar, Ahmad Muallimin, Fachrizal Ambia

Abstract

Background

Futsal is a high-intensity sport that requires optimal cardiorespiratory endurance to maintain performance throughout a match. VO₂ Max is a key indicator of aerobic capacity and overall physical fitness in athletes.

Objectives

This study aimed to evaluate the cardiovascular-respiratory endurance of SMAS Fatih Bilingual School futsal club players in Banda Aceh, Indonesia, and classify their VO₂ Max levels.

Methods

A quantitative descriptive-evaluative design was used, involving all 15 active futsal players through a total sampling technique. Cardiorespiratory endurance was assessed using the Harvard Step Test, and results were classified according to VO₂ Max norms for males aged 13–19 years. Data were analyzed descriptively to determine mean values, frequency distribution, and percentage classification.

Results

The mean VO₂ Max score was 62.78, categorized as “poor.” Distribution analysis showed that 33.33% of players were in the “very poor” category, 26.66% in “poor,” 26.66% in “moderate,” and 13.33% in “good.”

Conclusion

Most futsal players at SMAS Fatih Bilingual School demonstrated suboptimal aerobic fitness, which may hinder competitive performance. Implementing structured training programs targeting aerobic endurance is recommended to enhance performance and reduce fatigue during matches.

Keywords: Cardiorespiratory Endurance, VO₂ Max, Futsal, Harvard Step Test

Received: June, 10 2025. Accepted: June, 25 2025

*Correspondence: fachrulazmi@usk.ac.id

Fachrul Azmi

Correspondence Author Affiliate Universitas Syiah Kuala, Aceh, Indonesia.

Copyright: © 2025 by the authors. Published by KHATEC, Pontianak, Indonesia. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (Creative Commons Attribution-ShareAlike 4.0 International License), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



How to Cite: Azmi, F., Ifwandi, Masri, Zulfikar, Muallimin, A., & Ambia, F. (2025). Cardiorespiratory endurance assessment of high school futsal players: A case study at SMAS Fatih Bilingual School, Banda Aceh. *International Journal Emerging of Sport Science*, 1(2), 13-16

INTRODUCTION

Futsal has become one of the most widely played sports in Indonesia, attracting participants across diverse age groups and genders. While it is often dominated by male players, futsal has also gained popularity among women, reflecting its inclusive nature and accessibility (Halim & Indiarsa, 2016). As a variant of football, futsal is played indoors on a smaller pitch, with teams consisting of five players each. This format emphasizes fast-paced gameplay, technical precision, tactical adaptability, and high-intensity physical effort (Supriady, 2021).

To sustain optimal performance in such a demanding sport, players must possess various components of physical fitness, including endurance, speed, explosive power, agility, strength, and coordination. Among these, cardiorespiratory endurance—often measured through maximal oxygen uptake (VO₂ Max)—is particularly crucial. VO₂ Max refers to the maximum rate at which oxygen can be consumed during intense exercise, reflecting the combined efficiency of the cardiovascular, respiratory, and muscular systems (Nasution, 2014; Zakiyuddin, 2016). Higher VO₂ Max values are strongly associated with prolonged high-intensity performance, reduced fatigue, and improved match outcomes (Tamara & Nurrocmah, 2019).

Several factors influence VO₂ Max, including age, sex, genetics, nutrition, training regimen, and environmental conditions such as altitude (Indrayana & Yuliawan, 2019). Furthermore, the status of the cardiovascular, respiratory, and hematological systems directly affects oxygen delivery and utilization during exercise. Effective physical training programs—designed with systematic and progressive overload—are essential to improve aerobic fitness, enhance recovery rates, and maintain competitive performance levels (Yanti & Marisa, 2021).

Fatih Bilingual School, located in Banda Aceh, Indonesia, is known for its commitment to high-quality education, interactive learning, and extracurricular excellence. Its futsal team regularly competes in regional tournaments, such as the *Pascal Cup* (2023 and 2024), and often represents the school in provincial-level

competitions. However, observational assessments during friendly matches have revealed signs of performance decline during the later stages of play. These include reduced ball control, inaccurate passing and shooting, diminished off-ball movement, and slower responses during counterattacks—symptoms typically associated with inadequate cardiorespiratory endurance.

Despite the recognized importance of aerobic fitness in futsal, there is limited empirical research assessing VO₂ Max levels among high school players, particularly within Fatih Bilingual School. Understanding the players' aerobic capacity is essential for designing targeted conditioning programs that can enhance performance and reduce fatigue. Therefore, this study aims to evaluate the cardiorespiratory endurance of SMAS Fatih Bilingual School futsal players using the Harvard Step Test and classify their VO₂ Max levels according to established norms.

METHOD

Participant

The study involved all 15 active members of the SMAS Fatih Bilingual School futsal club in Banda Aceh, Indonesia. Participants were male students aged 16–18 years, all of whom regularly participated in the school's futsal training sessions and competitions. Due to the relatively small number of players, a total sampling technique was employed, ensuring that every team member was included in the research.

Research Design

This research employed a quantitative descriptive–evaluative design. Quantitative methods were used to obtain measurable data on the players' cardiorespiratory endurance, while the descriptive–evaluative approach aimed to profile their current fitness levels based on standardized assessment procedures (Sugiyono, 2012; Pratama & Nawawi, 2020). The Harvard Step Test served as the primary measurement tool, following standard protocols. Players stepped onto and off a bench at a set rhythm for a specified duration, after which heart rates were measured during recovery. Predicted VO₂ Max values were calculated and categorized according to norms for males aged 13–19 years (Parmar, 2013).

Data Analysis

Data were processed using descriptive statistics. The mean VO₂ Max score was calculated to determine the overall aerobic fitness classification of the team. Additionally, frequency distribution and percentage analysis were applied to classify players into four categories: *very poor*, *poor*, *moderate*, and *good*. These results were then interpreted to provide insights into the team's current endurance capacity and to inform recommendations for training improvements.

RESULTS AND DISCUSSION

Results

The results of the Harvard Step Test revealed the predicted VO₂ Max scores and corresponding fitness classifications of the 15 SMAS Fatih Bilingual School futsal players. The calculated mean VO₂ Max was **62.78**, which falls within the *poor* classification according to the VO₂ Max norms for males aged 13–19 years (Parmar, 2013).

VO₂ Max Classification

The frequency distribution showed that:

- a) **5 players (33.33%)** were classified as *very poor*
- b) **4 players (26.66%)** were classified as *poor*
- c) **4 players (26.66%)** were classified as *moderate*
- d) **2 players (13.33%)** were classified as *good*

This distribution indicates that more than half of the players (60%) fall into the *very poor* or *poor* categories, suggesting a general lack of optimal aerobic capacity within the team.

Table 1 presents the detailed VO₂ Max classification of the players.

Table 1. Predicted VO₂ Max classifications of SMAS Fatih Bilingual School futsal players

Classification	Frequency	Percentage (%)
Very Poor	5	33.33
Poor	4	26.66
Moderate	4	26.66
Good	2	13.33
Total	15	100

The high proportion of players in the lower fitness categories highlights the need for targeted aerobic conditioning programs. Players in the *moderate* and *good* categories generally demonstrated better performance consistency and quicker recovery during matches, whereas those in the *very poor* and *poor* categories exhibited earlier onset of fatigue, potentially compromising their tactical and technical execution.

Discussion

The results of this study indicate that the majority of SMAS Fatih Bilingual School futsal players have suboptimal cardiorespiratory endurance, as reflected by a mean VO₂ Max score of 62.78, which falls within the *poor* classification. More than half of the players (60%) were in either the *very poor* or *poor* categories, suggesting that their aerobic capacity is insufficient to sustain optimal performance throughout a full match.

Cardiorespiratory endurance plays a crucial role in futsal performance, as the sport is characterized by repeated high-intensity actions interspersed with brief recovery periods (Tamara & Nurrocmah, 2019). Players with low VO₂ Max levels are more likely to experience early fatigue, which can lead to reduced technical accuracy, slower decision-making, and diminished tactical discipline during later stages of play (Yanti & Marisa, 2021). This was consistent with observational reports during matches, where players displayed decreased ball control, inaccurate passing, and reduced off-ball movement after the midway point of the game.

Several factors may explain the low VO₂ Max values observed in this study. First, inconsistent participation in structured physical conditioning sessions was noted among some players, aligning with previous findings that training adherence significantly influences aerobic fitness development (Indrayana & Yuliawan, 2019). Second, lifestyle factors such as nutrition, rest, and recovery may not have been adequately optimized, which are known to affect cardiovascular efficiency (Zakiyuddin, 2016). Finally, differences in age and training experience could have contributed to the observed variation in VO₂ Max classifications, as younger and less experienced players often demonstrate lower endurance compared to senior players with longer training histories (Nasution, 2014).

The presence of players in the *moderate* and *good* categories suggests that improvement is achievable with appropriate interventions. Research has shown that targeted aerobic conditioning programs, including high-intensity interval training (HIIT) and small-sided games, can significantly improve VO₂ Max in futsal players within 6–8 weeks (Hadi, 2016). Additionally, integrating periodized training plans that combine aerobic, anaerobic, and strength components may further enhance endurance while supporting overall match performance.

From a practical standpoint, coaches at SMAS Fatih Bilingual School should consider implementing a structured preseason conditioning program, complemented by in-season maintenance sessions, to elevate the team's overall aerobic capacity. Emphasis should also be placed on player education regarding nutrition, hydration, and recovery strategies, as these factors can have cumulative effects on endurance and performance.

CONCLUSION

This study found that the cardiorespiratory endurance of SMAS Fatih Bilingual School futsal players was generally low, with a mean VO₂ Max score of 62.78 (*poor* category) and 60% of players falling into the *very poor* or *poor* classifications. Such fitness limitations likely contribute to early fatigue and reduced performance during matches. To address this, coaches should implement structured aerobic conditioning programs—such as high-intensity interval training and small-sided games—while encouraging consistent training attendance, proper nutrition, adequate recovery, and regular VO₂ Max monitoring. These measures are expected to improve endurance capacity and enhance team competitiveness in future competitions.

ACKNOWLEDGMENT

The author would like to express sincere gratitude to all those who contributed to the completion of this research. Special thanks are extended to the headmaster, teachers, and students of SMP Negeri 5 Kota Bima for their cooperation and participation throughout the study. The support and assistance provided during the data collection and training sessions were invaluable.

Appreciation is also extended to the supervisors and lecturers at the Physical Education Study Program, STKIP Taman Siswa Bima, for their continuous guidance, constructive feedback, and encouragement during the research process. Lastly, the author is deeply thankful to family and peers whose moral support and motivation made this study possible.

AUTHOR CONTRIBUTION STATEMENT

Fachrul Azmi⁸ conceptualized the study, designed the research methodology, and supervised the overall project. Ifwandi contributed to data collection and field testing. Masri assisted in data analysis and interpretation. Zulfikar contributed to literature review and drafting of the manuscript. Ahmad Muallimin participated in data collection and coordinate³ with the research site. Fachrizal Ambia provided critical revisions and final approval of the manuscript. All authors read and approved the final version of the article.

CONFLICT OF INTEREST AND FUNDING

There is no conflict of interest

REFERENCES

- Cheevers, K. (2007). *Harvard Step Test protocol*. Retrieved from <https://www.brianmac.co.uk/step.htm>
- Hadi, F. S. (2016). Pengaruh latihan ladder drills terhadap peningkatan kelincahan siswa U-17 di Persatuan Sepakbola Jajag Kabupaten Banyuwangi. *Jurnal Pendidikan Jasmani*, 26(1), 213–228.
- Halim, A., & Indiarso, D. (2016). [Details missing—please provide complete reference information for proper APA formatting].
- Indrayana, B., & Yuliawan, E. (2019). Penyuluhan pentingnya peningkatan VO₂ Max guna meningkatkan kondisi fisik pemain sepakbola Fortuna FC Kecamatan Rantau Rasau. *Jurnal Ilmiah Sport Coaching and Education*, 3(1), 41–50. <https://doi.org/10.21009/JSCE.03105>
- Nasution, S. (2014). [Details missing—please provide complete reference information for proper APA formatting].
- Parmar, P. (2013). *Norms for Harvard Step Test in adolescents*. [Details of publication].
- Pratama, T., & Nawawi, U. (2020). Tinjauan tingkat kondisi fisik atlet sepakbola Porma FC Sijunjung. *Jurnal Patriot*, 2, 549–564.
- Sugiyono. (2012). *Metode penelitian komunikasi: Kuantitatif, kualitatif, dan cara mudah menulis artikel pada jurnal internasional*. Alfabeta.
- Tamara, & Nurrocmah. (2019). Quality of sports physical fitness extracurricular participants. *Jurnal Sportif: Jurnal Penelitian Pembelajaran*, 5(2), 312–322.
- Yanti, E. D., & Marisa, D. (2021). Perbedaan VO₂ Max antara pemain bola basket dan pemain bola voli. *Jurnal Ilmiah Penjas*, 4(3), 233–240.
- Zakiyuddin. (2016). Analisis VO₂ Max pemain sepakbola usia 17–20 tahun di Club Bligo Putra Sidoarjo. *Jurnal Pendidikan Kepeleatihan Olahraga*, 1(1), 1–7.

Cardiorespiratory Endurance Assessment of High School Futsal Players: A Case Study at SMAS Fatih Bilingual School, Banda Aceh

ORIGINALITY REPORT

8%	%	8%	%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

- 1** Andi Nurcahyo, Faisal Yusuf, Diah Aryati Puji Lestari, Prima Januar Sastrawirya. "Zumba Exercise Training to Train Motor Movement & VO2 Max", Jurnal Pengabdian KOLABORATIF, 2025
Publication 2%
- 2** Justita Dura, Ditya Wardana. "The Sustainability Role of Women Entrepreneurs through the Digital Financial Literacy Movement", Population and Economics, 2024
Publication 1%
- 3** Argian Rizki Taufik, Amung Ma'mun, Mulyana Mulyana. "Dampak Shooting Three Point Plyometric dan Ladder terhadap Hasil Shooting Three Point Bola Basket", Gelanggang Olahraga: Jurnal Pendidikan Jasmani dan Olahraga (JPJO), 2020
Publication 1%
- 4** SCOTT R. SMALL, SHING CHAN, ROSEMARY WALMSLEY, LENNART VON FRITSCH et al. "Self-Supervised Machine Learning to 1%

Characterize Step Counts from Wrist-Worn Accelerometers in the UK Biobank", Medicine & Science in Sports & Exercise, 2024

Publication

-
- | | | |
|---|--|-----|
| 5 | Eka Iswahyudi, Muhammad Suhdy, Muhammad Supriyadi. "Perbandingan Daya Tahan Kardiorespirasi Antara Perokok Aktif dan Perokok Pasif Pemain Futsal Forza Lubuklinggau", e-SPORT: Jurnal Pendidikan Jasmani, Kesehatan dan Rekreasi, 2023 | 1% |
| 6 | Khoirul Fatihin, Gigih Siantoro, Muhammad Muhammad. "Kondisi Fisik Atlet Bola Voli Putra Puslatda Sebelum dan Saat Pandemi Covid 19", Gelanggang Olahraga: Jurnal Pendidikan Jasmani dan Olahraga (JPJO), 2021 | 1% |
| 7 | Didi Suryadi, Rubiyatno Rubiyatno. "Kebugaran jasmani pada siswa yang mengikuti ekstrakurikuler futsal", Jurnal Ilmu Keolahragaan, 2022 | 1% |
| 8 | Rohit Singh, Krishna Pal Sharma, Lalit Kumar Awasthi. "A machine learning-based ensemble model for securing the IoT network", Cluster Computing, 2024 | 1% |
| 9 | Yangyang Xie, Salman Alfarisi. "The Artistic Characteristics and Evolution of the Clown Dance in Gannan Tea-Picking Opera", Journal | <1% |
-

of Integrated Social Sciences and Humanities, 2025

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off